




Development of integrated livestock breeding and management strategies to improve animal health, product quality and performance in European organic and 'low input' milk, meat and egg production



Subproject 4: Laying Hens

**Ferry Leenstra, Veronika Maurer, Monique Bestman,
Esther Zeltner, Thea van Niekerk, Fabien Galea and
Berry Reuvekamp**



Wageningen, March 15

Low input systems for laying hens



Commercial production of eggs

- › Organic
- › Free range

**Hens receive complete diet (more or less ad libitum),
but have outside access**

**In poultry real low input is back yard farming (see FAO
E-conference on “Opportunities of the poultry breeding
programs for the family poultry production in
developing countries : The bird for the poor”)**



2

Goals



- › Develop a participatory system to test and optimize genotypes specific for free range and organic systems
- › Optimize management issues for free range and organic farms with special emphasis on diets and feather pecking
- › Analyze how the productive life of laying hens can be extended (consequences for health)
- › Analyze/optimize egg quality characteristics



3

Methods



Inventory among free range and organic farms in CH, F and NL

	Switzerland		France		Netherlands	
	Free range	Organic	Free range	Organic	Free range	Organic
N farms	35	91	32	11	48	57
Farm size	3.093	1.635	7.577	4.682	17.625	8.077
Egg production	244.1	241.9	247.0	245.4	244.9	231.0
Mortality (%)	5.9	6.6	4.9	4.7	6.6	12.0



4

Genotypes



>20 different genotypes:

- › 10 brands of brown hens (1 – 51 flocks/brand)
- › 3 brands of white hens (4-28 flocks/brand)
- › 4 brands of silver hens (3-15 flocks/brand)
- › 3 original genotypes
- › 73 mixed flocks (brown and white, brown and silver, white and silver)

Some, but not complete overlap between countries



5

Performance by genetic group



	White	Brown	Silver	Brown + Silver	Brown+ White	White+ Silver
Flocks	32	120	31	5	28	4
Free range (N eggs/hh)	248.7	246.2	237.8	248.0	200.0	NP
Organic (N eggs/hh)	243.5	239.1	227.2	254.3	240.8	243
Free range (%)	5.2	5.8	9.8	5.6	1.0	NP
Organic (%)	3.5	8	13.4	9.6	7.1	10.4

Brand and country to some extent confounded, but organic lower production, higher mortality
Silver more problems than White or Brown



6

Next flock different genotype?



	Same	Different	Don't know
Free range	103	25	12
Organic	66	97	8
Switzerland	69	77	0
France	32	6	0
The Netherlands	68	29	20

Free range less shift in genotype than organic
Switzerland: egg trader and/or hatchery decide



7

Workshops 'Ideal hen'



Workshops with farmers in CH and NL.

Results quite similar:

- › **Longevity (with or without moulting)**
- › **Adaptability (fast recovery)**
- › **Behaviour**
 - › **Curious, bold, calm, 'optimistic'**
 - › **Nesting behaviour, no smothering**
- › **Eating capacity, bit more body mass**
- › **Good persistence more important than high peak**



8

Next steps




Farm visits (20/system/country)

- › What determines slaughter age
- › More insight in management factors
- › Feather score, breast bone
- › Try new genotype?
- › Egg quality characteristics

- › Experimental setting
- › Genotype x diet interaction
 - › Genotype: new vs currently common? (or ...)
 - › Diet: with and without animal proteins



9



The authors gratefully acknowledge co-funding from the European Commission, under the Seventh Framework Programme for Research and Technological Development, for the Collaborative Project LowInputBreeds (Grant agreement No 222623)

The work of Wageningen LR and Louis Bolk is partly funded by the Dutch Ministry of Economics, Agriculture and Innovation



10