Simple soil carbon model for use in the HighCrop decision model

Jørgen E. Olesen, Aarhus University, Department of Agroecology 23 January 2014

The change in soil carbon (ΔC) (Mg C ha⁻¹ yr⁻¹) on an annual time scale is determined by the difference in net carbon additions (C_{net}) (Mg C ha⁻¹ yr⁻¹) and the degradation of native soil carbon (C_{deg}) (Mg C ha⁻¹ yr⁻¹).

$$\Delta C = C_{\text{net}} - C_{\text{deg}} \tag{1}$$

The amount of degradable soil C (C_{hum}) (Mg C ha⁻¹) is calculated from the amount of soil total nitrogen by assuming a C/N ratio of 11.

The degradation of soil C is calculated as a proportion of the native soil C using an annual degradation corresponding to 1.36 % estimated from long-term experiments at Askov research station in Denmark (Christensen, 1990):

$$C_{\rm deg} = 0.0136 \ C_{\rm hum}$$
 (2)

The net addition of carbon to the soil is calculated at the added amount (C_{add}) (Mg C ha⁻¹ yr⁻¹) multiplied by a humification coefficient (*h*):

$$C_{\rm net} = h \ C_{\rm add} \tag{3}$$

The humification coefficient is set to h = 0.15 for plant residues, corresponding to a 15% retention of added carbon over a number of years (Christensen, 2005). For manure the humification is set to 0.30, and this is increased to 0.40 for digested manure (Thomsen et al., 2012). For biochar the humification is set to 1.00.

For most crops the added carbon is calculated as

 $C_{\text{add}} = C_{\text{top}} + C_{\text{root}}$ (4) where C_{top} is added carbon in aboveground crop residues (Mg C ha⁻¹ yr⁻¹) and C_{root} is added carbon in belowground crop residues (Mg C ha⁻¹ yr⁻¹).

Sampling of roots of cereals at flowering have shown slightly higher root biomass in winter cereals compared with spring cereals, but little or no effect of fertilization and management (Chirinda et al., 2012). Therefore fixed values of added carbon in roots are used for each crop based on estimates used for the C-TOOL model (Taghizadeh-Toosi et al., 2013) (Table 1).

The added C in the top depends on dry matter crop yield (*Y*) (Mg DM ha⁻¹) and whether the straw is removed. It is assumed that dry matter has a carbon content of 45%:

$$C_{\rm top} = 0.45 \left[Y \left(1 - HI \right) \left(1 - f_{\rm straw} \right) \right]$$
(5)

where *HI* is the harvest index and f_{straw} is the proportion of above-ground residues (straw) that is removed when straw also is harvested. If straw is not harvested then f_{straw} is set to 0. The estimates of *HI* and *f*straw for the different crops are shown in Table 2.

Crop	$C_{\rm root}$ (Mg ha ⁻¹)
Winter cereal	1.6
Spring cereal	1.0
Winter rye	1.6
Spring oats	1.0
Pulse	1.1
Cereal-pulse intercrop	1.1
Lucerne	3.0
Grass-clover for cutting	4.0
Grass-clover for green manure	4.0
Grass-clover for grazing	4.0
Clover for seed	2.0
Sugar beet	0.6
Winter oilseed rape	1.6
Maize	1.5
Potato	0.6
Catch crop, poorly developed	0.7
Catch crop, medium developed	1.0
Catch crop, well developed	1.3

Table 1. Carbon additions in roots (C_{root}).

Table 2. Parameters for estimating carbon additions in tops of plants. *HI* is the harvest index and f_{straw} is the fraction of straw that is harvestable.

Сгор	HI	$f_{\rm straw}$
Winter cereal	0.45	0.45
Spring cereal	0.45	0.45
Winter rye	0.38	0.50
Spring oats	0.40	0.40
Pulse	0.42	0.40
Cereal-pulse intercrop (grain)	0.42	0.45
Lucerne for cutting	0.90	0.00
Grass-clover for cutting	0.95	0.00
Grass-clover for green manure	0.00	0.00
Grass-clover for grazing	0.80	0.00
Clover for seed	0.06	0.00
Sugar beet	0.70	0.50
Winter oilseed rape	0.37	0.50
Maize for silage	0.80	0.00
Potato	0.70	0.00
Catch crop, poorly developed	0.00	0.60
Catch crop, medium developed	0.00	0.70
Catch crop, well developed	0.00	0.80

For manure the amount of carbon added is calculated from the total N applied (N_{man}) (kg N ha⁻¹) in the manure using fixed C/N ratios (CN_{man}) for each manure type:

$$C_{\rm add} = CN_{man} N_{\rm man} / 1000$$

(2)

The C/N ratios for the different manure types are shown in Table 3.

Manure	CN _{man}
Cattle slurry	8.5
Pig slurry	4.3
Digested manure	5.0^{*}
Liquid manure (ajle)	2.0
Grass-clover silage	15.0
Deep litter, fresh	19.0
Deep litter, stored	13.5
Poultry manure, fresh	7.5
Poultry manure, stored	7.5
Compost	13.0

Table 3. C/N ratios of organic carbon to total N in various types of manure.

* Great variation between digested pig slurry (2.0) and digested cattle slurry/plant biomass (5-7.5)

References

- Chirinda, N., Olesen, J.E., Porter, J.R., 2012. Root carbon input in organic and inorganic fertilizerbased systems. Plant and Soil 359, 321-333.
- Christensen, B.T., 1990. Sædskiftets indflydelse på jordens indhold af organisk stof. II. Markforsøg på grov sandblandet lerjord (JB5), 1956-1985. Tidsskr. Planteavl 94, 161-169.
- Christensen, B.T., 2005. Kulstof i dyrket jord vurdering af potentiale for øget lagring. Side 103– 120 i DJF rapport Markbrug nr. 113. Drivhusgasser fra jordbruget – reduktionsmuligheder (J. E. Olesen, red.). Danmarks JordbrugsForskning, Tjele.
- Taghizadeh-Toosi, A., Hutchings, N.J., Vejlin, J., Christensen, B.T, Olesen, J.E., 2013. C-TOOL: A soil carbon model and its parameterisation. Ecological Modelling (in prep).
- Thomsen, I.K., Olesen, J.E., Møller, H.B., Sørensen, P., Christensen, B.T., 2012. Carbon dynamics and stabilization in soil after anaerobic digestion of dairy cattle feed and faeces. Soil Biology and Biochemistry 58, 82-87.